

Auslander-Reiten Quivers and the Coxeter complex

Shmuel ZELIKSON *

August 7, 2002

ABSTRACT : Let Q be a quiver of type ADE . We construct the corresponding Auslander-Reiten quiver as a topological complex inside the Coxeter complex associated with the underlying Dynkin diagram. We use the notion of chamber weights coming from the theory of the canonical basis of quantized enveloping algebras, and show this set has a special linearity property in our setting. Finally, we consider A_n case, and describe how Auslander-Reiten quivers correspond to particular wiring diagrams.

Introduction

Let D be a Dynkin diagram of type ADE , and Q a quiver obtained by orienting its edges. Associated with Q is a finite dimensional algebra over $\mathbb{C}$, the path algebra $\mathbb{C}Q$. The category of representations of the quiver Q over $\mathbb{C}$ identifies with the category $\text{mod } \mathbb{C}Q$ of finitely generated $\mathbb{C}Q$ modules. Gabriel's theorem states that isomorphism classes of indecomposables ("indecomposables") of this category are indexed by the positive roots of the root system Φ associated with D . Furthermore only quivers of ADE types have finitely many indecomposables.

The Auslander-Reiten quiver Γ_Q codifies the structure of the category $\text{mod } \mathbb{C}Q$. Vertices are the indecomposables, arrows are irreducible morphisms between them. Our aim is to construct Γ_Q as a topological complex $\tilde{\Gamma}_Q$, while retaining the spirit of Gabriel's theorem. We use the Coxeter complex Σ , that is the simplicial complex determined by hyperplanes perpendicular to the positive roots, in order to get a unified labelling of both vertices and arrows of $\tilde{\Gamma}_Q$.

The vertices of $\tilde{\Gamma}_Q$ are one dimensional rays inside Σ , that is weights lying in Weyl group orbits of the fundamental weights. To be more precise, Q defines a bilinear form R on the Euclidean space E generated by Φ which corresponds to the homological form at the level of $\text{mod } \mathbb{C}Q$. Every column of the matrix of R in the basis of simple roots defines a weight $-\rho_i$. The vertices of $\tilde{\Gamma}_Q$ obtain then out of the ρ_i 's by the same linear combinations which construct Φ^+ out of the simple roots $\alpha_1, \dots, \alpha_n$.

Q defines a particular reduced expression $\tilde{w}_0$ of the longest word of W , up to commutation of simple reflections, referred to as an adapted reduced expression. $\tilde{w}_0$ induces a total order $\mathcal{R}_{\tilde{w}_0}$ of Φ^+ , and the main result of [2] is that the pair $\tilde{w}_0, \mathcal{R}_{\tilde{w}_0}$

*AMS Mathematics Subject classification : 16G70, 17B10, 20F55.

encodes the structure of Γ_Q (Theorem 1.1) . Conversely, one can recover Γ_Q out of $\tilde{w}_0$ (Proposition 1.2).

Passing from the simple roots to fundamental weights, one associates with $\tilde{w}_0$ the set of chamber weights $\mathcal{C}_{\tilde{w}_0}$ (Definition 2.1). This set was introduced in [4] in the study of total positivity questions in real semi-simple groups. We show (Corollary 2.5) $\mathcal{C}_{\tilde{w}_0}$ is the set of vertices defined above using R . Actually one has linearity of the 1 : 1 correspondence $\varphi_{\tilde{w}_0} : \mathcal{R}_{\tilde{w}_0} \longrightarrow \mathcal{C}_{\tilde{w}_0}$, (Theorem 2.4), which turns out to be a restriction to Φ^+ of a linear application $-\varphi_R$ defined by R . We conjecture linearity is a characterization of reduced expressions $\tilde{w}_0$ adapted to quivers.

Two dimensional faces of Σ (planar cones) are uniquely determined by the two rays to which they are adjacent. They may therefore be seen as edges between vertices. This allows us to use the set of vertices $\mathcal{C}_{\tilde{w}_0}$ in order to associate with any reduced expression $\tilde{w}_0$ a quiver $\Gamma_{\tilde{w}_0}$ (Definition 3.5). This quiver depends on $\tilde{w}_0$ only up to commutation of simple reflections. When $\tilde{w}_0$ is adapted to a quiver Q , $\Gamma_{\tilde{w}_0}$ depends only on Q , so we can note it by $\tilde{\Gamma}_Q$. We show this quiver is isomorphic to Γ_Q (Theorem 3.7). We use a direct combinatorial approach, thus isomorphism is proved independently of linearity. Note however Theorem 2.4 defines naturally the isomorphism involved. The $\tilde{\Gamma}_Q$ being special cases of the $\Gamma_{\tilde{w}_0}$, we call the later abstract Auslander-Reiten quivers.

The last section describes as an example A_n type. This case is characterized by the fact all fundamental representations are minuscule. We can therefore index one-dimensional rays in Σ by Young columns, and arrows of $\Gamma_{\tilde{w}_0}$ become simply couples of column tableaux characterized by inclusion of indices (Lemma 4.1). This fact, combined with the existence of action of a Coxeter element in $\tilde{\Gamma}_Q$ (Proposition 2.7) similar to the well-known one on Γ_Q , allows us to give a combinatorial description (Proposition 4.3) of $\tilde{\Gamma}_Q$. On the theoretical level, use of Young columns makes the link between $\Gamma_{\tilde{w}_0}$ and the wiring diagram $\mathcal{WD}(\tilde{w}_0)$ of $\tilde{w}_0$, used in [3] as a key tool in the study of Lusztig's parameterizations of the canonical basis (consult the more recent [5] for use of chamber weights).

We are grateful to Bernhard Keller for helpful comments. We also wish to thank the referee for his remarks, which led us to improve the first version of this work, dealing only with A_n case. This work was partially supported by EC TMR network "Algebraic Lie Representations", grant no. ERB FMRX-CT97-01000.

1 Auslander-Reiten quivers

We recall in this section basic facts about Auslander-Reiten quivers, and proceed to study their link with reduced words of the longest word of the Weyl group.

Let us fix $\mathbb{C}$ as the ground field k . Consider a *ADE* Dynkin diagram D , of rank n , and index its vertices as in [14], page 53. For sake of simplicity we shall identify the set of simple roots $\{\alpha_1, \dots, \alpha_n\}$ with $I = \{1, \dots, n\}$.

We note by Φ the root system associated with D , E the real vector space in which it lies, $(,)$ the scalar product defined on E by the Cartan matrix, and by $\alpha_i^\vee$ the coroots. Let W be the Weyl group, $\{s_1, \dots, s_n\}$ the set of simple reflections, $l()$ the length function, and w_0 the unique longest element of W . w_0 induces an involution on

I (paragraphs (XI) of [7], Planches I,IV,V,VI,VII), which we shall note by $i \mapsto i^*$.

We obtain a quiver Q by orienting the edges of D . Q is given therefore as a couple (Q_0, Q_1) , with $Q_0 (= I)$ a set of vertices, and Q_1 a set arrows. In this work Q will always denote a fixed quiver whose underlying Dynkin diagram is of type ADE . Let $\text{mod } \mathbb{C}Q$ denote the category of finite type representations of Q over the complex numbers [9] 77.1. An object in this category is given by $M = (\underline{V}, \underline{\varphi})$, where with every $i \in I$ we associate a finite dimensional vector space V_i , and with every arrow $i \longrightarrow j$ in Q_1 a linear mapping $\varphi_{i,j} : V_i \longrightarrow V_j$. One obtains the simple objects by associating a one dimensional vector space to a vertex i , and 0 spaces to the others. They are therefore in 1 : 1 correspondence with the simple roots.

We shall note by $\text{Ind } Q$ the set of isomorphism classes $[M]$ of indecomposable representations in $\text{mod } \mathbb{C}Q$. The dimension vector of a representation M is the linear combination $\dim M = \sum_{i \in I} a_i \alpha_i$ with $a_i = \dim V_i$.

Gabriel's Theorem Let Q be a quiver of type ADE . Then the mapping $[M] \mapsto \dim M$ establishes a one to one correspondence between $\text{Ind } Q$ and Φ^+ . Furthermore, the quivers of type ADE are the only ones for which $\text{Ind } Q$ is a finite set.

The category $\text{mod } \mathbb{C}Q$ is hereditary. As observed by Ringel [18] (see also [19]), the homological form

$$\langle M, N \rangle := \dim_{\mathbb{C}} \text{Hom}(M, N) - \dim_{\mathbb{C}} \text{Ext}^1(M, N)$$

depends only on the dimension vectors. If we note by $[\beta]$ the isomorphism class of the indecomposable corresponding to $\beta \in \Phi^+$, then $\langle [\alpha], [\beta] \rangle = R(\alpha, \beta)$ where R is the bilinear form on E defined by

$$r_{i,j} := R(\alpha_i, \alpha_j) = \begin{cases} 1 & \text{if } i = j \\ -1 & \text{if } i \longrightarrow j \in Q_1 \\ 0 & \text{otherwise} \end{cases}$$

We shall refer to R as the Ringel form of Q


The structure of $\text{mod } \mathbb{C}Q$ is codified by the Auslander-Reiten quiver Γ_Q . Vertices are $\text{Ind } Q$, arrows are irreducible morphisms, namely morphisms that cannot be written as a non-trivial composition of two morphisms ([1] page 166). Γ_Q is endowed with translation τ ([1] page 225), which stratifies it into levels Γ_Q^i indexed by I . Such a level is of the form

$$(P_i(Q) =) \tau^{N_i-1} I_i(Q), \tau^{N_i-2} I_i(Q), \dots, \tau I_i(Q), I_i(Q)$$

with $P_i(Q)$ the projective cover of the simple module corresponding to $i \in I$, and $I_i(Q)$ the injective envelope of the simple module corresponding to $i^* \in I$.


A detailed construction of Γ_Q our case is given by [11] section 6.5. With D one associates an infinite quiver $\mathbb{Z}D$ [11] fig 13. page 49, whose vertices are $I \times \mathbb{Z}$. Arrows can exist between vertices only if their first coordinates are linked in D . The vertices of D are ordered in a natural way such that for i, j linked in D , the corresponding arrows

in $\mathbb{Z}D$ are given by the configuration



Thus $\mathbb{Z}D$ is an “infinite mesh” deduced out of D . One defines on it the Nakashima involution ν [11] page 48. The opposite quiver Q^{op} to Q may be realized inside $\mathbb{Z}D$ by a slice S_{proj} starting at $(1, 0)$. Its image by ν gives a slice S_{inj} which has the same arrows as Q . The Auslander-Reiten quiver then identifies with the subgraph of $\mathbb{Z}D$ delimited by S_{proj} to the left, and S_{inj} to the right. Vertices of S_{proj} (resp. S_{inj}) are the projectives in mod $\mathbb{C}Q$ (resp. injectives). Each translation level Γ_Q^i corresponds to the vertices in its realization inside $\mathbb{Z}D$ having i as first coordinate. One may deduce recursively the dimension vectors of the vertices of Γ_Q out of the slice S_{proj} by use of additivity ([11] page 50) (for examples, see [2] Examples 2.7, 2.8).

Example 1.1 : Consider the A_n type quiver $Q^{(d)} : \bullet \xrightarrow{1} \bullet \xrightarrow{2} \dots \xrightarrow{n} \bullet$, which is distinguished by the regularity of its structure. We reproduce the figure in [19] page 85, $\Gamma_{Q^{(d)}}$ is given by



where $M_{i,j+1} := [\alpha_i + \alpha_{i+1} + \dots + \alpha_j]$. We have $S_{proj} = \{M_{1,n+1}, M_{2,n+1} \dots M_{n,n+1}\}$, ($M_{n,n+1}$ corresponding to $(0, 1) \in \mathbb{Z}D$), and $S_{inj} = \{M_{1,n+1}, M_{1,n} \dots M_{1,3}, M_{1,2}\}$. For $i \in I$, the translation level $\Gamma_{Q^{(d)}}^i$ is i^{th} horizontal level from the top (i.e. containing $[M_{i,n+1}]$).

Given $w \in W$, we shall use the notation $\tilde{w} = s_{i_1} \dots s_{i_m}$ for a reduced expression of w , the $\sim$ indicating we are referring not to w but to its reduced expression. Recall one has for the longest element $l(w_0) = N$ where $N = \text{card} \Phi^+$. A fixed reduced expression $\tilde{w}_0 = s_{i_1} s_{i_2} \dots s_{i_N}$ induces a total ordering of the positive roots $\mathcal{R}_{\tilde{w}_0} = (\beta_1, \beta_2 \dots \beta_N)$ by

$$\beta_j = s_{i_1} s_{i_2} \dots s_{i_{j-1}}(\alpha_{i_j}).$$

$\mathcal{R}_{\tilde{w}_0}$ obtains as a linear refinement of convex partial order on Φ^+ . Convex (or normal) means a partial order $\preceq$ verifying conditions [3] :

$$\begin{aligned} & \forall \alpha, \beta \in \Phi^+, \text{ such that } \gamma = \alpha + \beta \in \Phi^+ : \\ & \text{either } \alpha \preceq \gamma \preceq \beta \text{ or } \beta \preceq \gamma \preceq \alpha. \end{aligned}$$

Conversely, a convex order $\preceq$, determines $\tilde{w}_0$, up to commutation of simple reflections, that is up to "2-moves" replacing $s_{i_j} s_{i_{j+1}}$ by $s_{i_{j+1}} s_{i_j}$, i_j and i_{j+1} not linked in D . If we consider two reduced expressions of w_0 as equivalent if one may be obtained from the other by a sequence of 2-moves, then the equivalence classes we obtain are known as commutation classes [2] 1.4. We have a 1 : 1 correspondence between them and convex orderings. All constructions in the sequel will actually depend only on the commutation class (or equivalently a convex ordering).

A quiver Q defines a convex ordering $\preceq_Q$ [19]. One may use the Ringel form, putting $\beta \preceq \gamma$ if $\langle \beta, \gamma \rangle > 0$, and taking the transitive closure of these relations. The same partial order may be obtained from Γ_Q , one has $\alpha \preceq \beta$ iff there is a path in Γ_Q from $[\alpha]$ to $[\beta]$. Equivalence of the two constructions follows from the fact arrows in Γ_Q are couples $[\alpha], [\beta]$ such that $R(\alpha, \beta) > 0$ and there is no γ such that $R(\alpha, \gamma) > 0$ and $R(\gamma, \beta) > 0$. We shall say $\preceq$ is the convex ordering adapted to Q , and use the same reference to reduced expressions in the corresponding commutation class.

We say, following [6] a vertex k is a sink of Q if there are only entering arrows into it. Call it a source if there are only exiting arrows out of it. If k is a sink of Q define $S_k Q$ as the quiver obtained by reversing the direction of the arrows with k as end, thus turning k into a source. A reduced expression $\tilde{w}_0 = s_{i_1} s_{i_2} \dots s_{i_N}$ is adapted to Q if and only if i_1 is a sink of Q , i_2 a sink of $S_{i_1} Q$, i_3 is a sink of $S_{i_2} S_{i_1} Q$ and so on.

Example 1.2 : Consider $Q = Q^{(d)} : \bullet \xrightarrow{1} \bullet \xrightarrow{2} \bullet$ of type A_3 . 1 is a source, 3 is a sink, and $Q' = S_3 Q$ is then

$$Q' : \bullet \xrightarrow{1} \bullet \xleftarrow{2} \bullet$$

We leave it to the reader to verify $\tilde{w}_0 = s_3 s_2 s_1 s_3 s_2 s_3$ is adapted to $Q^{(d)}$ while $\tilde{w}_0 = s_2 s_1 s_3 s_2 s_3 s_1$ is adapted to Q' . For arbitrary n , one may verify $\tilde{w}_0 = s_n s_{n-1} \dots s_1 s_n s_{n-1} \dots s_2 \dots s_n s_{n-1} s_n$ is adapted to the distinguished orientation Q_d .

Given Q , let us renumber its vertices in a compatible way with its arrows. Namely we renumber $\{\alpha_1, \dots, \alpha_n\}$ as $\{\beta_1, \dots, \beta_n\}$ such that $\beta_j \longrightarrow \beta_k$ in Q implies $j > k$. One then associates to Q ([6] §1.2) a Coxeter element $c = s_{\beta_n} \dots s_{\beta_2} s_{\beta_1}$. There might be several compatible indexations, yet by [6] Lemma 1.2, c is independent of a particular choice. For instance $c = s_3 s_2 s_1$ is the Coxeter element associated to the quiver $Q^{(d)}$ in the example above, and $c = s_1 s_3 s_2 (= s_3 s_1 s_2)$ the Coxeter element to the quiver Q' . Observe, as in the proof of [6] Theorem 1.2, $s_{\beta_n} \dots s_{\beta_2} s_{\beta_1}$ is a sequence adapted to Q . It is well known that translation τ of Γ_Q becomes action of the Coxeter element c at the level of dimension vectors.

If k is a sink of Q , and $Q' := S_k Q$ then the categories $\text{mod } \mathbb{C}Q$ and $\text{mod } \mathbb{C}Q'$ are closely linked by the BGP-reflection functor [9] 77.4 corresponding to s_k . Bedard [2] combines use of these functors together with the description of Γ_Q inside $\mathbb{Z}D$ in order to show how to obtain from Γ_Q all $\tilde{w}_0$ adapted to Q .

If for $i \in I$, the occurrences of s_i in $\tilde{w}_0$ are at positions $j_1, j_2 \dots j_r$ define $\mathcal{R}_{\tilde{w}_0}^i$ as the ordered r -tuple $(\gamma_1, \gamma_2, \dots, \gamma_r)$ with $\gamma_k = s_{i_1} s_{i_2} \dots s_{i_{j_k-1}}(\alpha_{j_k})$ Theorem 2.17 of [2] may

then be restated as follows :


Theorem 1.1 Let Q be a quiver of type ADE .

a) Associate with every vertex $[\beta]$ of Γ_Q the translation level i_β it belongs to. Read $\text{Ind } Q$ sequentially, in a manner compatible with the arrows (i.e $[\alpha] \longrightarrow [\beta]$ implies $[\alpha]$ appears before $[\beta]$). Replacing vertices $[\beta]$ by the corresponding i_β one produces an N -tuple $(i_1, i_2, \dots, i_N)$ giving a reduced expression adapted to Q . Furthermore, all $\tilde{w}_0$ adapted to Q are obtained this way.


b) Fix $\tilde{w}_0$ as in a). For all $i \in I$ taking dimension vectors establishes a $1 : 1$ correspondence (as ordered sets) between Γ_Q^i and $\mathcal{R}_{\tilde{w}_0}^i$.

Part b) tells us in particular each $[\alpha] \in \Gamma_Q^i$ has the form $\alpha = s_{i_1} s_{i_2} \dots s_{i_{j-1}}(\alpha_{i_j})$ with $i_j = i$. We shall note by $N_i(Q)$ the cardinality of Γ_Q^i (or of $\mathcal{R}_{\tilde{w}_0}^i$).

Example 1.3 : Consider the quiver $Q := \overset{1}{\bullet} \longrightarrow \overset{2}{\bullet} \longleftarrow \overset{3}{\bullet}$ of type A_3 . Γ_Q is



Replacing vertices by their translation levels one obtains



There are 4 ways of reading the vertices in a compatible way with the arrows $(2, 1, 3, 2, 1, 3)$, $(2, 1, 3, 2, 3, 1)$, $(2, 3, 1, 2, 1, 3)$, $(2, 3, 1, 2, 3, 1)$. The 4 corresponding reduced expressions form the commutation class determined by Q .

Theorem 1.1 allows us also to describe the opposite direction, recovering Γ_Q from a $\tilde{w}_0$ adapted to Q . Citing the referee, this result is well known to the experts, yet unpublished so we provide its proof.

Proposition 1.2 Let $\tilde{w}_0$ be a reduced expression adapted to a quiver Q of type ADE . Consider a couple of roots $\alpha = s_{i_1} s_{i_2} \dots s_{i_{j-1}}(\alpha_{i_j})$, $\beta = s_{i_1} s_{i_2} \dots s_{i_{k-1}}(\alpha_{i_k})$ in $\mathcal{R}_{\tilde{w}_0}$. $[\alpha] \longrightarrow [\beta]$ is an arrow in Γ_Q iff

- i) i_j and i_k are linked in the Dynkin diagram D ,
- ii) $j < k$,
- iii) For $j < p < k$, $i_p \neq i_j$.

Proof : Since Γ_Q lies inside $\mathbb{Z}D$, observe there are arrows only between translation levels whose indices are linked in D . We need therefore only to consider the case of i_j and i_k linked in D . Put $a := i_j$, $b := i_k$. As $\tilde{w}_0$ is adapted to Q , by Theorem 1.1 it obtains by reading translation levels of vertices of Γ_Q in a compatible manner with its arrows. Now the mesh structure of $\mathbb{Z}D$ implies an arrow from a vertex $[\alpha]$ on Γ_Q^a to a vertex $[\beta]$ on Γ_Q^b is immediately followed by an arrow from $[\beta]$ to the vertex $[\gamma]$ following $[\alpha]$ on the translation level Γ_Q^a , and vice-versa. In order to comply with this configuration, if we delete from $\tilde{w}_0$ all simple reflection s_{i_l} with $l \neq a, b$ we have to obtain a word either of the form $s_a s_b s_a s_b \dots$ or of the form $s_b s_a s_b s_a \dots$. By ii) and iii), our criterion gives then the same structure of arrows as the one of Γ_Q inside $\mathbb{Z}D$ $\square$.

Definition 1.3 Let $\tilde{w} = s_{i_1} \dots s_{i_m}$ be a reduced expression of $w \in W$. We shall say $\tilde{w}$ is alternating if for every couple j, k of indices linked in D , the word one obtains by erasing from $\tilde{w}$ all s_{i_r} , $i_r \neq j, k$ is either of the form $s_i s_j s_i \dots$ or of the form $s_j s_i s_j \dots$.

Lemma 1.4 Any reduced expression $\tilde{w}_0$ adapted to a quiver Q is alternating.

The proof is included in that of Proposition 1.2.

Example 1.4 : Consider $\tilde{w}_0 = s_3 s_2 s_1 s_3 s_2 s_3$. For the couple $\{1, 2\}$, we get by erasing occurrences of s_3 , the word $s_2 s_1 s_2$, and for the couple $\{2, 3\}$ erasing occurrences of s_1 gives $s_3 s_2 s_3 s_2 s_3$.

2 The linearity phenomenon

We import in this section the notion of chamber weights $\mathcal{C}_{\tilde{w}_0}$ from quantum groups and show in case $\tilde{w}_0$ is adapted to Q , the correspondence $\varphi_{\tilde{w}_0} : \mathcal{R}_{\tilde{w}_0} \longrightarrow \mathcal{C}_{\tilde{w}_0}$ is linear and given by the Ringel form.

Let us consider orbits of fundamental weights ω_i instead of positive roots.

Definition 2.1 Let W be a Weyl group of finite type, and let us fix a reduced expression $\tilde{w}_0$. The set of chamber weights associated with $\tilde{w}_0$ is the N -tuple $\mathcal{C}_{\tilde{w}_0} := (\mu_1, \mu_2, \dots, \mu_N)$ where $\mu_j = s_{i_1} s_{i_2} \dots s_{i_j}(\omega_{i_j})$.

Chamber weights were introduced in [4] as a key tool in the study of total positivity in semi-simple groups. Our convention is in opposite order to that in [4]. All elements of $\mathcal{C}_{\tilde{w}_0}$ are distinct ([4] 2.9). By its construction, $\mathcal{C}_{\tilde{w}_0}$ seen as a set depends only on the commutation class of $\tilde{w}_0$. Let us observe the cardinality of $\mathcal{F} = \bigcup_{i \in I} W \omega_i$ is greater than N , so unlike the positive roots case, $\tilde{w}_0$ actually “cuts out” a subset inside $\mathcal{F}$.

For any $\tilde{w}_0$, one has a natural 1 : 1 correspondence $\varphi_{\tilde{w}_0}$ between $\mathcal{R}_{\tilde{w}_0} = (\beta_1, \dots, \beta_N)$ and $\mathcal{C}_{\tilde{w}_0} = (\mu_1, \dots, \mu_N)$ defined by $\varphi_{\tilde{w}_0}(\beta_k) = \mu_k$, $k = 1, \dots, N$. Observe this bijection depends only on the commutation class, as interchange of two commuting reflections $s_{i_j}, s_{i_{j+1}}$ results in an interchange between β_j, β_{j+1} in $\mathcal{R}_{\tilde{w}_0}$, and μ_j, μ_{j+1} in $\mathcal{C}_{\tilde{w}_0}$.

If we consider columns of the matrix $(r_{i,j})$ of the Ringel form R as coordinate vectors in the basis of the ω_i 's, we obtain the following weights :

$$\forall i \in I : \rho_i := \sum_{l \in I} r_{l,i} \omega_l.$$

We note by φ_R the linear mapping from E to E defined by $\varphi_R(\alpha_i) = \rho_i$ for i in I .

Lemma 2.2 Suppose k is a sink of Q and $Q' := S_k Q$. Let R and R' be the corresponding Ringel forms. Then

$$s_k \varphi_{R'} = \varphi_R s_k$$

Proof : The lemma is a direct consequence of the behavior of Ringel forms under *BGP* reflection functors. Let $\{\rho_i\}_{i \in I}$, $\{\rho'_i\}_{i \in I}$ be the respective weights defined by R and R' . Note by D_k the set of $j \neq k$ linked to k in D .

The coefficient of ω_k in ρ_j verifies $[\rho_j : \omega_k] = 0$ for $j \neq k$, while $\rho_k = \omega_k - \sum_{j \in D_k} \omega_j$.

In comparison, we have for Q' :

$$\rho'_j = \begin{cases} \omega_k & j = k, \\ \rho_j - \omega_k & j \in D_k, \\ \rho_j & \text{otherwise.} \end{cases}$$

Let us show directly $\forall j \in I : \varphi_{R'}(\alpha_j) = s_k \varphi_R s_k(\alpha_j)$ (we give details for convenience of the reader).

Case 1 : $j = k$.

$$\begin{aligned} s_k \varphi_R s_k(\alpha_k) &= s_k(-\rho_k) \\ &= s_k(-\omega_k + \sum_{j \in D_k} \omega_j) \\ &= -\omega_k + \alpha_k + \sum_{j \in D_k} \omega_j. \end{aligned}$$

Now $\alpha_k = 2\omega_k - \sum_{j \in D_k} \omega_j$ so we get $s_k(-\rho_k) = \omega_k = \rho'_k = \varphi_{R'}(\alpha_k)$.

Case 2 : $j \in D_k$.

$$\begin{aligned} s_k \varphi_R s_k(\alpha_j) &= s_k \varphi_R(\alpha_j + \alpha_k) \\ &= s_k(\rho_j + \rho_k). \end{aligned}$$

$s_k(\rho_j) = \rho_j$ since $[\rho_j : \omega_k] = 0$. We have just seen $s_k(\rho_k) = -\omega_k$. We get therefore $s_k \varphi_R s_k(\alpha_j) = \rho_j - \omega_k = \rho'_j = \varphi_{R'}(\alpha_j)$

Case 3 : $j \neq k$ and not in D_k .

Immediate, $\rho_j = \rho'_j$ and s_k stabilizes both α_j and ρ_j $\square$

Corollary 2.3 Assume the expression $s_{i_1} s_{i_2} \dots s_{i_{k-1}}$ is adapted to Q . Note for $j = 1 \dots k-1 : Q_j := S_{i_{j-1}} S_{i_{j-2}} \dots S_{i_1} Q$ ($Q_1 := Q$) and R_j the Ringel form corresponding to Q_j . Then

$$\varphi_{R_1} s_{i_1} s_{i_2} \dots s_{i_{k-1}} = s_{i_1} s_{i_2} \dots s_{i_{k-1}} \varphi_{R_k}.$$

Proof : For any $j = 1, \dots, k-1$, i_j is a sink of Q_j so that $\varphi_{R_j} s_{i_j} = s_{i_j} \varphi_{R_{j+1}}$.

Theorem 2.4 (Linearity) Let $\tilde{w}_0$ a reduced expression adapted to Q of type *ADE*. Then the mapping $\varphi_{\tilde{w}_0}$ is linear and given by the Ringel form R :

$$\forall 1 \leq k \leq N, \mu_k = -\varphi_R(\beta_k)$$

Corollary 2.5 The $-\rho_i$'s, $i \in I$ are elements of $\mathcal{C}_{\tilde{w}_0}$, and $\mathcal{C}_{\tilde{w}_0}$ is obtained from them by the same linear combinations which construct Φ^+ out of the simple roots $\{\alpha_1, \dots, \alpha_n\}$.

Proof of Theorem 2.4 : Consider an arbitrary k , $1 \leq k \leq N$. By hypothesis, $\tilde{w}_0 = s_{i_1} s_{i_2} \dots s_{i_N}$ is adapted to Q so may apply Corollary 2.3 and its conventions (in particular $R_1 = R$).

$$\begin{aligned} \varphi_R(\beta_k) &= \varphi_{R_1} s_{i_1} s_{i_2} \dots s_{i_{k-1}}(\alpha_{i_k}) \\ &= s_{i_1} s_{i_2} \dots s_{i_{k-1}} \varphi_{R_k}(\alpha_{i_k}) \\ &= s_{i_1} s_{i_2} \dots s_{i_{k-1}} \varphi_{R_k}(s_{i_k}(-\alpha_{i_k})) \\ &= s_{i_1} s_{i_2} \dots s_{i_{k-1}} s_{i_k} \varphi_{R_{k+1}}(-\alpha_{i_k}) \\ &= -s_{i_1} s_{i_2} \dots s_{i_{k-1}} s_{i_k} \varphi_{R_{k+1}}(\alpha_{i_k}) \end{aligned}$$

Observe i_k is a source of Q_{k+1} . We have seen while proving Lemma 2.2, in that case $\varphi_{R_{k+1}}(\alpha_{i_k}) = \omega_{i_k}$, so finally

$$\varphi_R(\beta_k) = -s_{i_1} s_{i_2} \dots s_{i_k}(\omega_{i_k}) = -\mu_k. \quad \square$$

Examples 2.1 Consider A_3 type.

1) We have seen $\tilde{w}_0 = s_2 s_1 s_3 s_2 s_3 s_1$ is adapted to a quiver. $\mathcal{R}_{\tilde{w}_0}$ is given by

$$\begin{aligned} \beta_1 &= \alpha_2, & \beta_3 &= \alpha_2 + \alpha_3, & \beta_5 &= \alpha_1, \\ \beta_2 &= \alpha_1 + \alpha_2, & \beta_4 &= \alpha_1 + \alpha_2 + \alpha_3, & \beta_6 &= \alpha_3, \end{aligned}$$

and $\mathcal{C}_{\tilde{w}_0}$ by

$$\begin{aligned} \mu_1 &= \omega_1 - \omega_2 + \omega_3, & \mu_3 &= \omega_1 - \omega_2, & \mu_5 &= -\omega_1, \\ \mu_2 &= -\omega_2 + \omega_3, & \mu_4 &= -\omega_2, & \mu_6 &= -\omega_3. \end{aligned}$$

The matrix of the Ringel form is

$$\begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix}$$

μ_5, μ_1, μ_6 are the respective images by $\varphi_{\tilde{w}_0}$ of the simple roots, and indeed $-\rho_1 = -\omega_1 = \mu_5$, $-\rho_2 = \omega_1 - \omega_2 + \omega_3 = -\mu_1$, $-\rho_3 = -\omega_3 = \mu_6$. Finally, observe one has $\mu_2 = \varphi_{\tilde{w}_0}(\alpha_1 + \alpha_2)$, and as expected $-\rho_1 - \rho_2 = -\omega_2 + \omega_3$. Similarly, $\mu_3 = -\rho_2 - \rho_3$ and $\mu_4 = -\rho_1 - \rho_2 - \rho_3$.

2) The reduced expression $\tilde{w}_0 = s_2 s_1 s_2 s_3 s_2 s_1$ is not adapted to a quiver. $\mathcal{R}_{\tilde{w}_0} = (\beta'_1, \dots, \beta'_6)$ with

$$\begin{aligned} \beta'_1 &= \alpha_2, & \beta'_3 &= \alpha_1, & \beta'_5 &= \alpha_2 + \alpha_3, \\ \beta'_2 &= \alpha_1 + \alpha_2, & \beta'_4 &= \alpha_1 + \alpha_2 + \alpha_3, & \beta'_6 &= \alpha_3, \end{aligned}$$

while $\mathcal{C}_{\tilde{w}_0} = (\mu'_1, \dots, \mu'_6)$ with

$$\begin{aligned} \mu'_1 &= \omega_1 - \omega_2 + \omega_3, & \mu'_3 &= -\omega_1 + \omega_3, & \mu'_5 &= -\omega_2, \\ \mu'_2 &= -\omega_2 + \omega_3, & \mu'_4 &= -\omega_1, & \mu'_6 &= -\omega_3. \end{aligned}$$

In this case, linearity fails. Observe $\mu'_2 = \varphi_{\tilde{w}_0}(\alpha_1 + \alpha_2) = -\omega_2 + \omega_3$ while $\mu'_3 = \varphi_{\tilde{w}_0}(\alpha_1)$, $\mu'_1 = \varphi_{\tilde{w}_0}(\alpha_2)$ verify $\mu'_3 + \mu'_1 = -\omega_2 + 2\omega_3$.

The set $\mathcal{A}$ of adapted convex orderings (commutation classes) is a small subset of $\mathcal{O}$ of all convex orderings, we provide the table below for small ranks :

	A_3	A_4	A_5	A_6	D_4	D_5
card $\mathcal{A}$	4	8	16	32	8	16
card $\mathcal{O}$	8	62	908	24968	182	13198

Conjecture $\tilde{w}_0$ is adapted to a quiver iff the mapping $\varphi_{\tilde{w}_0}$ is linear.

Computer testing shows linearity fails for all convex orderings in $\mathcal{O} \setminus \mathcal{A}$ for the Dynkin types in the table above. It is not clear to us at the moment how linearity should be modified in order to describe $\varphi_{\tilde{w}_0}$ for arbitrary commutation classes.

The set $\mathcal{C}_{\tilde{w}_0}$ may be stratified into levels the same way $\mathcal{R}_{\tilde{w}_0}$ is, namely the i^{th} level $\mathcal{C}_{\tilde{w}_0}^i$ is the set of weights $\mu_k = s_{i_1} \dots s_{i_k}(\omega_{i_k})$ with i_k equal to i

Lemma 2.6 For all $i \in I$, the last element of $\mathcal{C}_{\tilde{w}_0}^i$ is $-\omega_{i^*}$.

Proof : Consider the last appearance $s_{j_k} = s_i$ in $\tilde{w}_0$. All following reflections s_{j_m} with $m > k$ stabilize ω_i . Hence $s_{j_1} s_{j_2} \dots s_{j_k} \omega_i = w_0 \omega_i = -\omega_{i^*}$ $\square$

Proposition 2.7 Let $\tilde{w}_0$ be adapted to Q , and c the Coxeter element associated to Q . Then for all $i \in I$:

$$\mathcal{C}_{\tilde{w}_0}^i = (c^{N_i-1}(-\omega_{i^*}), c^{N_i-2}(-\omega_{i^*}), \dots, c(-\omega_{i^*}), -\omega_{i^*})$$

where $N_i = N_i(Q)$.

Proof : Recall $c = s_{j_1} s_{j_2} \dots s_{j_n}$ with j_1 a sink of $Q_1 := Q$, j_2 a sink of $Q_2 := S_1 Q$ and so on. As $(j_1, j_2, \dots, j_n)$ is a permutation of $\{1, \dots, n\}$, every arrow in Q is inverted twice in the sequence $S_{j_n} \dots S_{j_1} Q$ so have $Q = Q_{n+1}$. Using Corollary 2.3 and $R_{n+1} = R_1 = R$, we get $c\varphi_R = \varphi_R c$. By Theorem 2.4 we can replace φ_R by $\varphi_{\tilde{w}_0}$ so that :

$$(*) \quad c\varphi_{\tilde{w}_0} = \varphi_{\tilde{w}_0} c.$$

We already know by Auslander-Reiten theory the levels of $\mathcal{R}_{\tilde{w}_0}$ are given by the action of c . It remains to apply $(*)$ to get the statement of the proposition. $\square$

Remark : If $\tilde{w}_0$ is not adapted to a quiver Q , the levels $\mathcal{C}_{\tilde{w}_0}^i$ may not be given through the action of a Coxeter element. We do not have for the moment a Weyl group model for the levels in general case.

3 The Coxeter complex construction

In this section we consider elements of $\mathcal{C}_{\tilde{w}_0}$ as one dimensional rays in the Coxeter complex Σ . The structure of Σ allows us to construct arrows, and associate a quiver $\Gamma_{\tilde{w}_0}$ with $\tilde{w}_0$. When $\tilde{w}_0$ is adapted to a quiver Q , we show we recover Γ_Q .

Recall the geometry of the root system Φ is captured by the Coxeter complex Σ (see [8] chapters 1-3, [7] chapter V). Geometrically Σ is the simplicial complex induced by the hyperplanes orthogonal to the positive roots. Maximal elements are the open chambers. Recall a parabolic subgroup of W is a subgroup $W^J = \langle s_j \rangle_{j \in J}$ with J a subset of I . Σ can be constructed out of W as the complex of all left classes of all parabolic subgroups partially ordered by inverse inclusion.

It would be more convenient for us to work in Σ^{op} , the inversely ordered complex, as the partial order becomes plain inclusion. Note $W^{(J)} := \langle s_j \rangle_{j \in I \setminus J}$. The maximal

elements (facets) of Σ^{op} are left classes of maximal parabolics $W^{(i)}$ (equivalently, one dimensional rays). Two facets $wW^{(i)}$, $w'W^{(j)}$ of Σ^{op} are adjacent iff their intersection is non-empty, which in that case, it is a left class of $W^{(i,j)}$.

Let us recall now basic facts about parabolic subgroups, one may consult for details [13] Chapter 1 or [12] Chapters 1 and 2 (Geck and Pfeiffer work with right classes, but all the arguments can be easily adapted to left classes).

Fix $J \subset I$. W^J is itself a Coxeter group, and as such is endowed with its proper length function l_J .

Proposition 3.1 ([12] 1.2.10) Consider $w \in W^J$. Then $l_J(w) = l(w)$. Moreover, if $\tilde{w} = s_{i_1} \dots s_{i_m}$ is any reduced expression of w inside W , then $\forall k$, i_k is in J .

Proposition 3.2 ([12] 2.1.1) Define

$$X_J = \{w \in W \mid l(ws) > l(w) \forall s \in J\}.$$

a) Every left class wW^J contains a unique element $u \in X_J$ and u is the unique element of minimal length inside wW^J .

b) For any element $y \in W^J$, if we write $y = ux$ with $x \in W^J$, then $l(y) = l(u) + l(x)$.

Consider $J \subset K \subset I$, and note by X_J^K the set of minimal representatives of the left classes of W^J in W^K

Lemma 3.3 ([12] 2.2.5)

$$X_J = X_K X_J^K.$$

Finally, $W^{(i)}$ is the stabilizer of ω_i , thus a weight $\mu = w\omega_i$ may be identified with the left class $wW^{(i)}$. If u is the minimal length element inside $wW^{(i)}$ we shall say $u\omega_i$ is the minimal presentation of μ .

Lemma 3.4 ([4] 2.7) Let $\mu = w\omega_i$, and $\tilde{w} = s_{i_1} s_{i_2} \dots s_{i_m}$ a reduced expression of w . The minimal representation of μ is obtained by erasing from $\tilde{w}$ all factors i_k for which $\langle \omega_i, \beta_k^\vee \rangle = 0$, where $\beta_k = s_{i_m} s_{i_{m-1}} \dots s_{i_{k+1}}(\alpha_{i_k})$.

Chamber weights can be seen as facets of Σ^{op} by identification above. We shall feel free to consider them as appropriate either as weights or left classes.


Definition 3.5

Let $\tilde{w}_0$ be a reduced expression of w_0 in W a Weyl group of finite type. The abstract Auslander-Reiten quiver $\Gamma_{\tilde{w}_0}$ associated with $\tilde{w}_0$ is defined as follows

- The set of vertices is the totally ordered set $\mathcal{C}_{\tilde{w}_0} = (\mu_1, \dots, \mu_N)$, seen as lying inside Σ^{op} .
- There is an arrow from $\mu_j = w_j W^{(i_j)}$ to $\mu_k = w_k W^{(i_k)}$ iff
 - i) i_j and i_k are linked in Dynkin diagram;
 - ii) $w_j W^{(i_j)}$ and $w_k W^{(i_k)}$ are adjacent in Σ ;
 - iii) $j < k$.

We are forced to introduce *iii*) as intersection of left classes is non-oriented. Observe also that by considering only "Dynkin couples" in *i*), $\Gamma_{\tilde{w}_0}$ actually depends only on the commutation class $[\tilde{w}_0]$ of $\tilde{w}_0$. Definition 3.5 provides therefore a unique quiver $\tilde{\Gamma}_Q$ associated to the reduced expressions adapted to a quiver Q . Our aim is to show it is isomorphic to Γ_Q . We shall follow a combinatorial approach, as suggested by the referee.

Example 3.1 : Consider the quiver $Q : \bullet^1 \longrightarrow \bullet^2 \longleftarrow \bullet^3$ of type A_3 . An adapted expression to Q is $\tilde{w}_0 = s_{i_1} s_{i_2} \dots s_{i_6}$ with $\underline{i} = (2, 1, 3, 2, 3, 1)$. $\Gamma_{\tilde{w}_0}$ is



where $w_k = s_{i_1} \dots s_{i_k}$, $k = 1, \dots, 6$.

Note for example

$$\begin{aligned} w_1 W^{(2)} &= \{s_2, s_2 s_1, s_2 s_3, s_2 s_1 s_3\} \\ w_2 W^{(1)} &= \{s_2 s_1, s_2 s_1 s_2, s_2 s_1 s_3, s_2 s_1 s_2 s_3, s_2 s_1 s_3 s_2, s_2 s_1 s_2 s_3 s_2\} \end{aligned}$$

are adjacent since $w_1 W^{(2)} \cap w_2 W^{(1)} = \{s_2 s_1, s_2 s_1 s_3\} (= s_2 s_1 W^{(1,2)})$, while $w_1 W^{(2)}$ and $w_6 W^{(1)}$ are not since $w_1 W^{(2)} \cap w_6 W^{(1)} = \emptyset$. We leave it to the reader to verify that there is an arrow between vertices on adjacent levels (=Dynkin couples) exactly when intersection of left classes is non empty.

Let $\tilde{w} = s_{i_1} s_{i_2} \dots s_{i_m}$ a reduced expression of an arbitrary element $w \in W$, $w \neq e$. We can define a partial set of chamber weights $\mathcal{C}_{\tilde{w}} = (\mu_1, \dots, \mu_n)$ with $\mu_j = s_{i_1} s_{i_2} \dots s_{i_{j-1}}(\mu_{i_j})$, and a partial quiver $\Gamma_{\tilde{w}}$ as above.

Proposition 3.6 Assume $\tilde{w}$ is alternating. Then combinatorial the description of Proposition 1.2 is valid for $\Gamma_{\tilde{w}}$. There is an arrow $\mu_j \longrightarrow \mu_k$ if and only if

- i) i_j and i_k are linked in the Dynkin diagram D ;
- ii) $j < k$;
- iii) For $j < p < k$, $i_p \neq i_j$.

Proof : We proceed by induction on $l(w)$. If $l(w) = 1$ there is nothing to prove. Suppose the proposition is valid for an alternating reduced expression $\tilde{w} = s_{i_1} \dots s_{i_m}$, and consider an alternating $\tilde{w}' = s_r \tilde{w}$ with $l(w') = l(w) + 1$. Note $\mathcal{C}_{\tilde{w}'} = (\mu'_1, \dots, \mu'_{m+1})$.

For any $x, y \in W$, $x W^{(i)} \cap y W^{(j)} \neq \emptyset \iff s_r x W^{(i)} \cap s_r y W^{(j)} \neq \emptyset$. Using induction hypothesis for $\tilde{w}$, the combinatorial description is valid for arrows $\mu'_j \longrightarrow \mu'_k$ inside $\Gamma_{\tilde{w}'}$ with $j \geq 2$. It remains to study arrows out of μ'_1 , that is intersections $s_r W^{(r)} \cap s_r s_{i_1} \dots s_{i_k} W^{(i_k)}$. By the above, this amounts to consider intersections $W^{(r)} \cap s_{i_1} \dots s_{i_k} W^{(i_k)}$. An arrow $\mu'_1 \longrightarrow \mu'_{k+1}$ exists in $\Gamma_{\tilde{w}'}$ iff such an intersection is non-empty. We shall consider $1 \leq k \leq m$ with i_k linked to r in D and put $w_k = s_{i_1} \dots s_{i_k}$.

Let us show first all arrows $\mu'_1 \longrightarrow \mu'_{k+1}$ given by the criterion appear in $\Gamma_{\tilde{w}'}$. If k verifies for all $1 < p < k$, $i_p \neq r$, as also $i_k \neq r$, we have $w_k \in W^{(r)}$. Hence $W^{(r)} \cap w_k W^{(i_k)} \neq \emptyset$ as expected.

Conversely, let us show there are no more arrows $\mu'_1 \longrightarrow \mu'_{k+1}$ in $\Gamma_{\tilde{w}'}$ than those given by the criterion. We suppose therefore there is p , $1 < p < k$ such that $i_p = r$. We can take p maximal with that property. $\tilde{w}$ is alternating by hypothesis, hence $i_{p+1}, i_{p+2} \dots i_{k-1}$ are different from r and i_k . By definition of the simple reflections we have the following coefficients ($\alpha_{i_p} = \alpha_r$) :

$$(*) \begin{cases} [s_{i_{k-1}} s_{i_{k-2}} \dots s_{i_{p+1}}(\alpha_{i_p}) : \alpha_r] = 1 \\ [s_{i_{k-1}} s_{i_{k-2}} \dots s_{i_{p+1}}(\alpha_{i_p}) : \alpha_{i_k}] = 0 \end{cases}$$

Now $s_{i_k}(\alpha_{i_p}) = \alpha_r + \alpha_{i_k}$, so we obtain from (*) :

$$[s_{i_k} s_{i_{k-1}} \dots s_{i_{p+1}}(\alpha_{i_p}) : \alpha_{i_k}] > 0$$

and therefore $(\omega_{i_k}, s_{i_k} s_{i_{k-1}} \dots s_{i_{p+1}}(\alpha_{i_p}^\vee)) \neq 0$. By Lemma 3.4, $s_{i_p} = s_r$ appears in the reduced expression $\tilde{u}$ for minimal representative u of $w_k W^{i_k}$ obtained by erasing procedure out of $\tilde{w}_k$.

Consider a left class of $W^{(r, i_k)}$ inside $w_{i_k} W^{(i_k)}$. By Lemma 3.3, its minimal representative is of the form $y = ux$ with x a minimal representative of a class of $W^{(r, i_k)}$ inside $W^{(i_k)}$. Take any reduced expression $\tilde{x}$ of x . $\tilde{y} = \tilde{u}\tilde{x}$ is a reduced expression of y , in which s_r appears. By Proposition 3.1 this means $y \notin W^{(r)}$, hence $yW^{(r, i_k)}$ is not included in $W^{(r)}$. We see no left class of $W^{(r, i_k)}$ can be contained in both $W^{(r)}$ and $w_k W^{(i_k)}$, so there is no arrow in $\Gamma_{\tilde{w}'}$ from μ'_1 towards $\mu'_{i_{k+1}}$ $\square$

Remark : Experimental data suggests the statement of the proposition is actually true in general, without the assumption that $\tilde{w}$ is alternating (see example 3.2 below). For this reason we insist that the third condition in the combinatorial characterization be $i_p \neq i_j$ while $i_p \neq i_j$, i_k would have been sufficient for the adapted case.

Theorem 3.7 (Isomorphism)


Let Q be a quiver of type ADE , and R its Ringel form. Then φ_R induces isomorphism of oriented graphs of Γ_Q onto $\tilde{\Gamma}_Q$.

Proof : Let $\tilde{w}_0$ be a reduced expression adapted to Q . Combining Theorems 1.1 and 2.4, we get a correspondence between vertices of Γ_Q and $\tilde{\Gamma}_Q$ given by φ_R . By Lemma 1.4, $\tilde{w}_0$ is alternating, hence applying Propositions 1.2 and 3.6, φ_R extends to an isomorphism of oriented graphs.

Theorem 3.7 justifies the use of the term Abstract Auslander-Reiten quiver for $\Gamma_{\tilde{w}_0}$. There is a finite dimensional algebra, behind the commutation class corresponding to Q , namely $\mathbb{C}Q$. Definition 3.5 arises the question of associating a finite dimensional algebra with any commutation class. One aim would be controlling the corresponding parameterizations of the canonical basis (see [17] for some adapted cases). Another application of finite dimensional algebra techniques in general reduced expressions setting, might be solving the problem of realizability of the set of commutation classes with braid moves (see [2]) as a 1-skeleton of a polytopal complex. This problem is also known as realizability of second Bruhat order [20], we point for instance to [10] as

representing the current trend of research concerning this problem. We observe for the moment, even the number of commutation classes in A_n case is unknown.

Example 3.2 : We have seen linearity fails for non-adapted cases. The same applies for the “mesh” structure coming from $\mathbb{Z}D$. Take $\tilde{w}_0 = s_2 s_1 s_2 s_3 s_2 s_1$. One may check $s_2 s_1 W^{(1)} \cap s_2 s_1 s_2 s_3 s_2 W^{(2)} = s_2 s_1 s_3 s_2 W^{(1,2)}$. $\Gamma_{\tilde{w}_0}$ is given by



We see there are two arrows from $s_2 s_1 W^{(1)}$ towards the second level. Observe also $\tilde{w}_0$ is no longer alternating, erasing s_3 we get $s_2 s_1 s_2 s_2 s_1$. The combinatorial description of Proposition 3.6 still applies.

4 The A_n case : wiring diagrams

We describe in this section A_n case. We use Young tableaux (columns) to describe $\tilde{\Gamma}_Q$. On the theoretical level, we describe briefly the correspondence of abstract Auslander-Reiten quivers with wiring diagrams used in [3].

Linearity allows us to compute easily the vertices of $\Gamma_{\tilde{w}_0}$, however, we are unable for the moment to provide an generalized adjacency criteria inside Σ^{op} , which would give in a straight-forward manner the arrows. There are in the literature case by case combinatorial descriptions of Σ according to Dynkin type, we restrict ourselves here to the simplest case : A_n .

All fundamental representations $E(\omega_i)$ of a simple complex lie algebra of type A_n are minuscule, so that $W\omega_i$ is the set of weights of $E(\omega_i)$. Furthermore as $E(\omega_i)$ is isomorphic to $\bigwedge^i E(\omega_1)$, it has a basis of weight vectors indexed by all Young columns of size i , filled in a strictly increasing manner with indices in $\{1, \dots, n+1\}$. We get therefore an identification of $W\omega_i$ (equivalently the set of left classes of $W^{(i)}$) with multi-indices $J = (j_1, j_2 \dots j_i)$, $1 \leq j_1 < j_2 < \dots < j_i \leq n$. Conversely, given a strictly increasing multi-index J of cardinality i (that is a Young column of size i), one may recover its weight as the sum of weights of its boxes

$$wt(J) = \sum_{k=1}^i wt(\boxed{j_k}),$$

with $wt(\boxed{j_k}) = \omega_k - \omega_{k-1}$, as such a box corresponds to a weight vector inside $E(\omega_1)$ (by convention $\omega_0 = \omega_{n+1} = 0$).

In A_n case, W is the symmetric group $\mathcal{S}_{n+1}$, so that simple reflections are the transpositions $(i \ i+1)$, and action on weight orbits becomes usual $\mathcal{S}_{n+1}$ action on sets of indices. Let us note for two integers $a < b$ the interval $(a, a+1, \dots, b)$ by $[a, b]$.

Proposition 4.1 Fix i between 1 and n , and consider two left cosets $wW^{(i)}$, $w'W^{(i+1)}$, with J , J' corresponding multi-indices. Then $wW^{(i)}$ and $w'W^{(i+1)}$ are adjacent in Σ^{op} if and only if $J \subset J'$.

Proof : The closure of the dominant chamber is a fundamental domain for the action of W . Furthermore pairs $(W^{(i)}, w'W^{(i+1)})$ correspond to left classes of $W^{(i,i+1)}$ inside $W^{(i)}$ which form a single $W^{(i)}$ orbit. Hence, the adjacent pairs $(wW^{(i)}, w'W^{(i+1)})$ in Σ^{op} form one orbit under diagonal action, that of $(W^{(i)}, W^{(i+1)})$.

The multi-indices corresponding to $W^{(i)}$ and $W^{(i+1)}$ are $[1, i]$ and $[1, i+1]$. As Weyl group action preserves inclusion of multi-indices we get from the discussion above that $wW^{(i)}$, $w'W^{(i+1)}$ adjacent implies $J \subset J'$.

For the converse we need to show any couple (J, J') with $J \subset J'$ can be obtained by diagonal action from $([1, i], [1, i+1])$. Since all multi-indices of cardinality i form a single Weyl group orbit, it is enough to consider couples of the form $([1, i], J')$. Since $[1, i] \subset J'$, we must have $J' = [1, i] \cup \{k\}$ with $k \geq i+1$. If $k > i+1$, then $s_{i+1}s_{i+2}\dots s_{k-1}J' = [1, i+1]$ while $s_{i+1}s_{i+2}\dots s_{k-1}$ stabilizes $[1, i]$. Thus

$$([1, i], J') = s_{k-1}s_{k-2}\dots s_{i+1}([1, i], [1, i+1]) \quad \square$$

Let us now describe $\tilde{\Gamma}_Q$ in terms of column tableaux. It seems somewhat simpler to use the classical method of Coxeter translation, instead of linearity.

As $W \cong \mathcal{S}_{n+1}$, the Coxeter element c defined by Q is a $n+1$ cycle. Let us fix a cycle writing $[c] = (a_1 a_2 \dots a_{n+1})$ by imposing $a_{n+1} = n+1$. In order to work inside $[c]$, we shall use $\overline{m}$ for $m \bmod n+1$. Finally let $[c]_{\overline{m}, i}$ be the multi-index obtained by ordering the "interval" $\{a_{\overline{m}}, a_{\overline{m}+1}, \dots, a_{\overline{m}+i-1}\}$.

Lemma 4.2 Define for $i \in I$, $\overline{m}_i$ recursively by $\overline{m}_1 = n+1$ and for $i \geq 2$

$$\overline{m}_i = \begin{cases} \overline{m}_{i-1} & \text{if } \bullet \xrightarrow{i^*} \bullet^{i^*+1} \text{ in } Q \\ \overline{m}_{i-1} - 1 & \text{if } \bullet \xleftarrow{i^*} \bullet^{i^*+1} \text{ in } Q \end{cases}$$

Then $[c]_{\overline{m}_i, i}$ is the multi-index $(i^*+1, i^*+2, \dots, n+1)$.

Proof : We proceed by induction on n . For case $n=2$ there is nothing to prove. Consider Q of type A_n , and let Q' be its sub-quiver with vertices 1 to $n-1$, with c' the corresponding Coxeter element, and m' defined for c' as above.

Case 1 : Q is obtained from Q' by adding $\bullet^{n-1} \xrightarrow{n} \bullet$. n is a sink, hence $c = s_n c' = (nn+1)c'$. If $c' = (b \dots a n)$ then $c = (b \dots a n+1 n)$. We have therefore $\overline{m}_2 = \overline{m}_1$.

Case 2 : Q is obtained from Q' by adding $\bullet^{n-1} \xleftarrow{n} \bullet$. n is a source, hence $c = c' s_n = c'(nn+1)$. If $c' = (b \dots a n)$ then $c = (b \dots a n n+1)$. We get $\overline{m}_2 = \overline{m}_1 - 1$.

In both cases, the cycle writing of c is obtained from that of c' by replacing n by either $nn+1$ or $n+1n$. The validity of the Lemma for c' in cases $i=2 \dots n-1$ implies its validity for c in cases $i=3 \dots n$. Also, the recursive formula for $\overline{m}_i$ for $i \leq 3$ is the same as that for $\overline{m}'_{i-1}$ $\square$

We see c has a "segment property" : for any j , indices $j, j+1, \dots, n+1$ form a "connected" band (modulo $n+1$) in the cycle writing of c . This property leads to the description of levels of $\tilde{\Gamma}_Q$. Let us note by $J_j^{(i)}(Q)$ the multi-index of the j^{th} vertex from the right (starting with 0) on the i^{th} level of $\tilde{\Gamma}_Q$.

Proposition 4.3 Consider Q and fix $i \in I$, then

a) ([2] Corollary 2.20 (a)) The cardinality $N_i(Q)$ of the i^{th} level of $\tilde{\Gamma}_Q$ is $(n + 1 + a_i - b_i)/2$ where a_i (respectively b_i) is the number of arrows in the unoriented path from i to i^* that are directed towards i (respectively i^*).

b) For $j = 0, 1, \dots, N_i(Q) - 1$ one has

$$J_j^{(i)}(Q) = [c]_{\overline{m_i - j}, i}$$


Proof of b) : Fix $i \in I$. By Lemma 2.6, the first weight from the right on the i^{th} level of $\tilde{\Gamma}_Q$ is $-\omega_{i^*}$. It corresponds to the multi-index $(i^* + 1, i^* + 2, \dots, n + 1)$. By the Lemma above, the proposition is true for $j = 0$.

For $1 \leq j \leq N_i(Q) - 1$, we use Proposition 2.7 :

$$\begin{aligned} J_j^{(i)}(Q) &= c^{-j} J_0^i(Q) \\ &= c^{-j} [c] \\ &= [c]_{\overline{m_i - j}, i} \square \end{aligned}$$

We see getting $\tilde{\Gamma}_Q$ out of Q requires two operations : computing the cycle writing of the Coxeter element associated to Q , and the cardinalities $N_i(Q)$. Both involve working directly with Q . Once vertices are known, getting the arrows is immediate due to Lemma 4.1.

Example 4.1 : Consider the quiver $Q : \overset{1}{\bullet} \xrightarrow{\quad} \overset{2}{\bullet} \xleftarrow{\quad} \overset{3}{\bullet}$ of type A_3 . $1^* = 3$ so that $a_1 = 1, b_1 = 1$ hence $N_1(Q) = 2$. We obtain in a similar manner $N_2(Q) = 2, N_3(Q) = 2$. The Coxeter element c is the cycle (2134) . The levels of $\tilde{\Gamma}_Q$ are given respectively by $\tilde{\Gamma}_Q^1 : (\cdot \cdot 3 \cdot), (\cdot \cdot \cdot 4)$; $\tilde{\Gamma}_Q^2 : (\cdot 13 \cdot), (\cdot \cdot 34)$; $\tilde{\Gamma}_Q^3 : (\cdot 134), (2 \cdot 34)$. Considering inclusion between sets of indices we get the arrows :




Reduced expressions $\tilde{w}_0$ of the longest word in W play a central role in the theory of the canonical basis [15] [16] of the positive part $U_q(\mathfrak{n}^+)$ of the quantized enveloping algebra $U_q(\mathfrak{g})$. As this base is difficult to compute, Lusztig's model consists in approximating it by a basis composed of PBW monomials. Every convex ordering of Φ^+ (or commutation class) gives rise to a particular PBW base, which provides an indexation of the canonical basis. The control over these parameterizations involves a geometrical interpretation of $\tilde{w}_0$, the wiring diagram $\mathcal{WD}(\tilde{w}_0)$. We refer to [3] §2.3 for the

details, and reduce the presentation here to example 4.2. Our conventions are “opposite” (twisted by $*$) compared to those of [3].

Given $\tilde{w}_0 = s_{i_1} s_{i_2} \dots s_{i_N}$ in $W \cong \mathcal{S}_{n+1}$, the wiring diagram $\mathcal{WD}(\tilde{w}_0)$ is an arrangement of $n + 1$ pseudo-lines of the plane $\mathbb{R}^2$, going from $-\infty$ to $+\infty$. The pseudo-lines cross pairwise in points $x_1, \dots, x_N$, situated in levels (top to bottom) in accordance with $\tilde{w}_0$, namely the crossing x_k is on the level i_k . We get a braid reversing the order of the lines, and which partitions $\mathbb{R}^2$ into zones (or chambers). One numbers the pseudo-lines from 1 to $n + 1$ as in the example (at x coordinate $-\infty$: from top to bottom). This gives a labelling of each zone Z by the set $J(Z)$ of indices of lines passing above it. Each zone has a distinct label. Observe [3] geometrically $\mathcal{WD}(\tilde{w}_0)$ depends only on the commutation class of $\tilde{w}_0$, and there is a 1 : 1 bijection between the set of commutation classes of A_n and possible wiring diagrams with $n + 1$ strands.

Example 4.2 : Take $\tilde{w}_0 = s_2 s_1 s_3 s_2 s_3 s_1$ (type A_3). $\mathcal{WD}(\tilde{w}_0)$ is :



Consider the set of zones bounded to the left by a crossing point. These are in bijection with crossing points x_k . Let us note therefore by Z_k the zone whose leftmost point is x_k in $\mathcal{WD}(\tilde{w}_0)$, and associate to it the level of x_k .

Lemma 4.4

a) The label $J(Z_k)$ is $w_k \cdot (1, 2, \dots, i_k)$ where $w_k = s_{i_1} s_{i_2} \dots s_{i_k}$.

b) Two chambers Z_j on the i^{th} level and Z_k on the $(i + 1)^{\text{th}}$ level are adjacent if and only if a pseudo-line L_m passes between them, that is if and only if $J(Z_k) = J(Z_j) \cup \{m\}$.

Proof : Pass a vertical line p_k through each crossing x_k . This divides the plane into vertical bands $\mathcal{V}_0, \mathcal{V}_1, \dots, \mathcal{V}_N$. $\mathcal{V}_0$ lies between $-\infty$ and p_1 , for $k = 1, \dots, N - 1$, $\mathcal{V}_k$ lies between p_k and p_{k+1} , and $\mathcal{V}_N$ lies between p_N and $+\infty$.

Note by J_k the ordered set of indices of pseudo-lines passing through $\mathcal{V}_k$, read top to bottom. Initially $J_0 = (1, 2, \dots, n)$. If we note $i = i_k$, the crossing x_k consists in interchanging the lines in positions i and $i + 1$. Suppose $J_{k-1} = (w_{k-1}(1), w_{k-1}(2), \dots, w_{k-1}(i), w_{k-1}(i + 1), \dots, w_{k-1}(n))$, then

$$\begin{aligned} J_k &= (w_{k-1}(1), w_{k-1}(2), \dots, w_{k-1}(i + 1), w_{k-1}(i), \dots, w_{k-1}(n)) \\ &= (w_{k-1} s_i(1), w_{k-1} s_i(2), \dots, w_{k-1} s_i(i), w_{k-1} s_i(i + 1), \dots, w_{k-1} s_i(n)) \\ &= (w_k(1), w_k(2) \dots w_k(n)). \end{aligned}$$


Proceeding by induction we get for $k = 1 \dots N$, $J_k = (w_k(1), w_k(2), \dots, w_k(n))$. Part a) follows then by observing for a zone Z_k , the label $J(Z_k)$ consists in the first i_k indices of J_k .

Part b) follows directly from our definition of labelling of the zones in $\mathcal{WD}(\tilde{w}_0)$ $\square$

We may consider labels of zones Z_k as Young columns. Lemma 4.4 a) shows this establishes a bijection with $\mathcal{C}_{\tilde{w}_0}$ (thus the term chamber weights used in [4]). Comparing Lemma 4.1 with Lemma 4.4 b), we see arrows in $\Gamma_{\tilde{w}_0}$ correspond to adjacencies of zones in $\mathcal{WD}(\tilde{w}_0)$. We invite the reader to compare Examples 4.1 and 4.2. Observe the non-left bounded zones may be considered as trivial, as they appear in any wiring diagram, and always have the same label. We therefore have a natural 1 : 1 correspondence between the set of abstract Auslander-Reiten quivers and that of wiring diagrams.

Remark : Wiring diagrams were generalized to B_n, C_n cases in [4]. No such construction exists in D_n or E_n cases.

In A_n case, we may summarize our results in the following figure :



We obtain another instance of the link between theories of finite dimensional algebras and of the canonical basis of $U_q(\mathfrak{g})$.

References

- [1] AUSLANDER, M.; REITEN, I.; SMALO, S.O.; *Representation theory of Artin algebras*; Cambridge Studies in Advanced Mathematics. 36, Cambridge University Press, (1995).
- [2] BEDARD, R.; *On commutation classes of reduced words in Weyl groups*; Eur. J. Comb. **20** (1999), 483-505.
- [3] BERENSTEIN, A.; FOMIN, S.; ZELEVINSKY, A.; *Parameterizations of canonical bases and totally positive matrices*; Adv. Math. **122** (1996), 49-149.
- [4] BERENSTEIN, A.; ZELEVINSKY, A.; *Total positivity in Schubert varieties*; Comm. Math. Helv. **72** (1997), 128-156.
- [5] BERENSTEIN, A.; ZELEVINSKY, A.; *Tensor product multiplicities, canonical bases, and totally positive varieties*; Invent. Math. **143** (2001), 77-128.
- [6] BERNSTEIN, I.N.; GELFAND, I.M.; PONOMAREV, V.A.; *Coxeter functors and Gabriel's theorem*; Russ. Math. Surveys **28** (1973), 17-32.

- [7] BOURBAKI, N.; *Groupes et algèbres de Lie. Chapitres 4, 5 et 6*; Elements de Mathématiques Masson, Paris (1981).
- [8] BROWN, K.S.; *Buildings* Springer-Verlag New-York (1989).
- [9] CURTIS, C.W.; REINER, I.; *Methods of representation theory with applications to finite groups and orders Vol. II.*; Wiley New-York (1987).
- [10] FELSNER, S.; ZIEGLER, G.; *Zonotopes associated with higher Bruhat orders*; Discrete Math. **241** (2001), 301-312.
- [11] GABRIEL, P.; *Auslander-Reiten sequences and representation-finite algebras*; in Representation theory I, Proc. Workshop, Ottawa 1979, Springer Lect. Notes Math. 831, 1-71.
- [12] GECK, M.; PFEIFFER, G.; *Characters of finite Coxeter groups and Iwahori-Hecke algebras* London Mathematical Society Monographs. New Series. 21. Oxford: Clarendon Press (2000).
- [13] HUMPHREYS, J.E.; *Reflection groups and Coxeter groups* Cambridge studies in advanced mathematics **29**, Cambridge University Press (1990).
- [14] KAC, V.G.; *Infinite dimensional Lie algebras. 3rd ed*; Cambridge University Press (1990).
- [15] KASHIWARA, M.; *On crystal bases of the q -analogue of universal enveloping algebras*; Duke Math. J. **63** (1991), 465-516.
- [16] LUSZTIG, G.; *Canonical bases arising from quantized enveloping algebras*; J. Am. Math. Soc. **3** (1990), 447-498.
- [17] REINEKE, M.; *On the coloured graph structure of Lusztig's canonical basis*; Math. Ann. **307** (1997) 705-723.
- [18] RINGEL, C.M.; *Representations of K -species and bimodules*; J. Alg. **41** (1976), 269-302.
- [19] RINGEL, C.M.; *PBW-bases of quantum groups*; J. Reine Angew. Math. **470** (1996), 51-88.
- [20] ZIEGLER, G.; *Higher Bruhat orders and cyclic hyperplane arrangements*; Topology **32** (1993) 259-279.

S. Zelikson, Laboratoire S.D.A.D., unité CNRS FRE 2272
 Département de Mathématiques
 Université de Caen B.P. 5186
 14032 Caen Cedex FRANCE
 E-mail : Shmuel.Zelikson@math.unicaen.fr